

Flokkar G þvagfæralyf og kyuhormónalyf; H hormónalyf; M vöðvasjúkdóma og beinagrindarlyf

<i>Classes of medications</i>	<i>No use %</i>	<i>Constant use %</i>	<i>Temporary use %</i>	<i>Use as needed %</i>	<i>Gender difference P value</i>	<i>Relative risk (95%CI)</i>
G. Genitourinary and sex hormones						
Female hormones women	77,8	12,9	9,2			
Drug used for benign prostatic hypertrophy men	77,4	16,2	6,4			
Drugs used for urinary frequency and incontinence	89,0	3,3	7,7		0,26	1,18 (0,87-1,61)
Karlar	89,3	2,6	9,0			1
Konur	89,0	3,6	7,0			
H. Systemic Hormonal preparations						
Glucocorticoids	97,2	1,2	2,6		0,038	
Karlar	95,8	1,6	1,0			1,02 (0,57-1,83)
Konur	98,0	1,0	1,6			1
Thyroid preparations	89,1	8,9	2,0		0,02	0,64 (0,46-0,91)
Karlar	92,0	6,0	2,0			1
Konur	87,6	10,5	2,0			
M. Musculoskeletal system						
Nonsteroidal anti-inflammatory agents	86,9	2,8	8,6	1,6	0,56	
Karlar	87,4	3,0	8,6	1,0		0,94 (0,71-1,25)
Konur	86,6	2,6	8,8	2,0		1
Coxibs	86,3	0,6	13,1	0,1	0,33	
Karlar	88,4	0,6	11,0	0		0,78 (0,59-1,04)
Konur	85,1	0,6	14,2	0,1		1
Bisphosphonates	89,1	4,2	6,7		0,000	
Karlar	95,0	1,4	3,6			0,35 (0,23-0,54)
Konur	85,9	5,7	8,4			1
Raloxifene						
konur	95,8	2,0	2,2			

Table III b. Class G: genitourinary and sex hormones; H: Systemic hormonal preparations; M: Musculoskeletal system. Proportional use %.