

Table I. Results of testing for antibodies against *Borrelia burgdorferi sensu latu* in serum samples from patients in Iceland from 2011 to 2015, broken down by year and antibody type. Number and percentage, n (%).

	Total	2011	2012	2013	2014	2015
N	552	85	112	116	124	115
CSF	73 (13.2)	8 (9.4)	16 (14.3)	17 (14.7)	19 (15.3)	13 (11.3)
<hr/>						
ELISA						
IgM	549 (99.5)	84 (98.8)	111 (99.1)	116 (100)	123 (99.2)	115 (100)
Positive	180 (32.6)	34 (40)	39 (34.8)	47 (40.5)	33 (26.6)	27 (23.5)
Negative	369 (66.8)	50 (58.8)	72 (64.3)	69 (59.5)	90 (72.6)	88 (76.5)
IgG	549 (99.5)	84 (98.8)	111 (99.1)	116 (100)	123 (99.2)	115 (100)
Positive	102 (18.5)	53 (62.4)	35 (31.3)	14 (12.1)	6 (4.8)	16 (13.9)
Negative	447 (81)	31 (36.5)	76 (67.9)	102 (87.9)	117 (94.4)	99 (86.1)
<hr/>						
Western blot						
IgM	186 (33.7)	38 (44.7)	38 (33.9)	46 (39.7)	33 (26.6)	31 (27)
Positive	22 (4)	4 (4.7)	3 (2.7)	6 (5.2)	7 (5.6)	2 (1.7)
Indeterminable	95 (17.2)	29 (34.1)	10 (8.9)	17 (14.7)	16 (12.9)	23 (20)
Negative	69 (12.5)	5 (5.9)	25 (22.3)	23 (19.8)	10 (8.1)	6 (5.2)
IgG	112 (20.3)	34 (40)	35 (31.3)	15 (12.9)	7 (5.6)	21 (18.3)
Positive	14 (2.5)	2 (2.4)	3 (2.7)	5 (4.3)	0 (0)	4 (3.5)
Indeterminable	54 (9.8)	28 (32.9)	4 (3.6)	1 (0.9)	7 (5.6)	14 (12.2)
Negative	44 (8)	4 (4.7)	28 (25)	9 (7.8)	0 (0)	3 (2.6)

Abbreviations: CSF - cerebrospinal fluid, ELISA - enzyme linked immunosorbent assay.