



Figure 1. Number of patients (y-axis) diagnosed with PSC for every year (x-axis) from 1992 to 2012.

<i>Authors, year</i>	Patients (n)	Incidence/100.000	Male ratio	Age at diagnosis (years)	IBD	Country
<i>Boberg et al. 1998¹¹</i>	17	1,3	x%	x	x	Norway
<i>Bambha et al. 2003⁶</i>	22	0,9	68%	40 (m)	73%	USA
<i>Kingham et al. 2004⁷</i>	53	0,9	62%	52 (M)	62%	UK
<i>aplan et al. 2007⁸</i>	49	0.9	55%	41 (M)	74%	Canada
<i>Card et al. 2008⁹</i>	149	0,4	64%	55 (m)	48%	UK
<i>Lindkvist et al, 2010¹⁰</i>	199	1,2	71%	39 (M)	76%	Sweden
<i>Guðnason et al, 2018</i>	42	0,7	67%	34 (M)	88%	Iceland

x=missing values, m=mean, M=median

Table I. Incidence, sex, age, IBD in PSC patients in previous studies.

Bloodtests	Patients (n)	Value at diagnosis (M)	Interquartile range (IQR)	Reference values
ALP	28	256 U/L	143-638 U/L	35-105 U/L
ALAT	30	118 U/L	66-221 U/L	KK 10-70 U/L Kvk 10-45 U/L
ASAT	31	73 U/L	50-108 U/L	KK 10-45 U/L Kvk 10-35 U/L
Gamma-GT	31	310 U/L	198-485 U/L	KK <115 U/L Kvk <75 U/L
Bilirubin	24	29 µmol/L	9-51 µmol/L	5-25 µmol/L
Albumin	18	38 g/L	35-43 g/L	36-45 g/L
Creatinine	26	71 µmol/L	60-80 µmol/L	50-100 µmol/L

Table II. Blodtest results at diagnosis.